

FINAL JEE-MAIN EXAMINATION – APRIL, 2024


(Held On Saturday 06th April, 2024)

TIME : 3 : 00 PM to 6 : 00 PM

CHEMISTRY

TEST PAPER WITH SOLUTION


SECTION-A



The **correct** arrangement for decreasing order of electrophilic substitution for above compounds

- (1) (IV) > (I) > (II) > (III)
 (2) (III) > (I) > (II) > (IV)
 (3) (II) > (IV) > (III) > (I)
 (4) (III) > (IV) > (II) > (I)

Ans. (2)



62. Molality (m) of 3 M aqueous solution of NaCl is:
 (Given : Density of solution = 1.25 g mL⁻¹, Molar mass in g mol⁻¹ : Na-23, Cl-35.5)

- (1) 2.90 m (2) 2.79 m
 (3) 1.90 m (4) 3.85 m

Ans. (2)

Sol. 3 moles are present in 1 litre solution

$$\text{molality} = \frac{3 \times 1000}{1.25 \times 1000 - [3 \times 58.5]} = 2.79 \text{ m}$$

63. The incorrect statements regarding enzymes are:


- (A) Enzymes are biocatalysts.
 (B) Enzymes are non-specific and can catalyse different kinds of reactions.
 (C) Most Enzymes are globular proteins.
 (D) Enzyme - oxidase catalyses the hydrolysis of maltose into glucose.

Choose the correct answer from the option given below:


- (1) (B) and (C) (2) (B), (C) and (D)
 (3) (B) and (D) (4) (A), (B) and (C)

Ans. (3)

Sol. Direct NCERT Based




Consider the above chemical reaction. Product "A" is:



Ans. (2)

Sol.




65. During the detection of acidic radical present in a salt, a student gets a pale yellow precipitate soluble with difficulty in NH₄OH solution when sodium carbonate extract was first acidified with dil. HNO₃ and then AgNO₃ solution was added. This indicates presence of:

- (1) Br⁻ (2) CO₃²⁻
 (3) I⁻ (4) Cl⁻

Ans. (1)

Download the new ALLEN app & enroll for Online Programs [CLICK HERE TO DOWNLOAD](#)



66. How can an electrochemical cell be converted into an electrolytic cell ?

- (1) Applying an external opposite potential greater than E_{cell}^0
- (2) Reversing the flow of ions in salt bridge.
- (3) Applying an external opposite potential lower than E_{cell}^0 .
- (4) Exchanging the electrodes at anode and cathode.

Ans. (1)

Sol. Applied external potential should be greater than E_{cell}^0 in opposite direction.

67. Arrange the following elements in the increasing order of number of unpaired electrons in it.


- (A) Sc (B) Cr
 (C) V (D) Ti
 (E) Mn

Choose the correct answer from the options given below:

- (1) (C) < (E) < (B) < (A) < (D)
- (2) (B) < (C) < (D) < (E) < (A)
- (3) (A) < (D) < (C) < (B) < (E)
- (4) (A) < (D) < (C) < (E) < (B)

Ans. (4)

Sol. Unpaired electron



68. Match List-I with List-II.

List-I	List-II
Alkali Metal	Emission Wavelength in nm
(A) Li	(I) 589.2
(B) Na	(II) 455.5
(C) Rb	(III) 670.8
(D) Cs	(IV) 780.0


Choose the correct answer from the options given below:

- (1) (A)-(I), (B)-(IV), (C)-(III), (D)-(II)
- (2) (A)-(III), (B)-(I), (C)-(IV), (D)-(II)
- (3) (A)-(IV), (B)-(II), (C)-(I), (D)-(III)
- (4) (A)-(II), (B)-(IV), (C)-(III), (D)-(I)





Ans. (2)

Sol. Fact Based



69. The major products formed:



A and B respectively are:




Ans. (2)



70. The incorrect statement regarding the geometrical isomers of 2-butene is:

- (1) cis-2-butene and trans-2-butene are not interconvertible at room temperature.
- (2) cis-2-butene has less dipole moment than trans-2-butene.
- (3) trans-2-butene is more stable than cis-2-butene.
- (4) cis-2-butene and trans-2-butene are stereoisomers.

Ans. (2)



Cis-but-2-ene has higher Dipole moment than trans-but-2-ene.

71. Given below are two statements:

Statement I: PF_5 and BrF_5 both exhibit sp^3d hybridisation.

Statement II: Both SF_6 and $[\text{Co}(\text{NH}_3)_6]^{3+}$ exhibit sp^3d^2 hybridisation.

In the light of the above statements, choose the correct answer from the options given below:


- (1) **Statement I** is true but **Statement II** is false
- (2) Both **Statement I** and **Statement II** are true
- (3) Both **Statement I** and **Statement II** are false
- (4) **Statement I** is false but **Statement II** is true

Ans. (3)



Both Statement (1) and (2) are false.

72. The number of ions from the following that are expected to behave as oxidising agent is:





- (1) 3
- (2) 4
- (3) 1
- (4) 2


Ans. (4)

Sol. Due to inert pair effect; Tl^{+3} and Pb^{+4} can behave as oxidising agents.

73. Identify the product (A) in the following reaction.



Ans. (2)



74. The correct statements among the following, for a "chromatography" purification method is:

- (1) Organic compounds run faster than solvent in the thin layer chromatographic plate.
- (2) Non-polar compounds are retained at top and polar compounds come down in column chromatography.
- (3) R_f of a polar compound is smaller than that of a non-polar compound.
- (4) R_f is an integral value.

Ans. (3)

Sol. Non polar compounds are having higher value of R_f than polar compound.



Download the new ALLEN app
& enroll for Online Programs

CLICK HERE TO
DOWNLOAD

75. Evaluate the following statements related to group 14 elements for their correctness.
- (A) Covalent radius decreases down the group from C to Pb in a regular manner.
- (B) Electronegativity decreases from C to Pb down the group gradually.
- (C) Maximum covalence of C is 4 whereas other elements can expand their covalence due to presence of d orbitals.
- (D) Heavier elements do not form $p\pi-p\pi$ bonds.
- (E) Carbon can exhibit negative oxidation states.
- Choose the **correct** answer from the options given below:

- (1) (C), (D) and (E) Only (2) (A) and (B) Only
(3) (A), (B) and (C) Only (4) (C) and (D) Only

Ans. (1)

- Sol.** (A) Down the group; radius increases
(B) EN does not decrease gradually from C to Pb.
(C) Correct.
(D) Correct.
(E) Range of oxidation state of carbon ; -4 to +4

76. Match List-I with the List-II

List-I
Reaction

- (A) $N_{2(g)} + O_{2(g)} \rightarrow 2NO_{(g)}$
(B) $2Pb(NO_3)_{2(s)} \rightarrow 2PbO_{(s)} + 4NO_{2(g)} + O_{2(g)}$
(C) $2Na_{(s)} + 2H_2O_{(l)} \rightarrow 2NaOH_{(aq)} + H_{2(g)}$
(D) $2NO_{2(g)} + 2OH_{(aq)}^- \rightarrow NO_{2(aq)}^- + NO_{3(aq)}^- + H_2O_{(l)}$

List-II
Type of redox reaction

- (I) Decomposition
(II) Displacement
(III) Disproportionation
(IV) Combination






Choose the **correct** answer from the options given below:

- (1) (A)-(I), (B)-(II), (C)-(III), (D)-(IV)
(2) (A)-(III), (B)-(II), (C)-(I), (D)-(IV)
(3) (A)-(II), (B)-(III), (C)-(IV), (D)-(I)
(4) (A)-(IV), (B)-(I), (C)-(II), (D)-(III)


Ans. (4)

- Sol.** A \rightarrow (IV)
B \rightarrow (I)
C \rightarrow (II)
D \rightarrow (III)

77. Consider the given reaction, identify the major product P.



Ans. (4)



78. The correct IUPAC name of $[PtBr_2(PMe_3)_2]$ is:

- (1) bis(trimethylphosphine)dibromoplatinum(II)
(2) bis[bromo(trimethylphosphine)]platinum(II)
(3) dibromobis(trimethylphosphine)platinum(II)
(4) dibromodi(trimethylphosphine)platinum(II)

Ans. (3)


Sol. Dibromo bis(trimethylphosphine) platinum (II)



Download the new **ALLEN** app
& enroll for **Online Programs**

CLICK HERE TO
DOWNLOAD

83. When 'x' × 10⁻² mL methanol (molar mass = 32 g; density = 0.792 g/cm³) is added to 100 mL water (density = 1 g/cm³), the following diagram is obtained.



x = (nearest integer)

[Given: Molal freezing point depression constant of water at 273.15 K is 1.86 K kg mol⁻¹]

Ans. (543)


Sol. $\Delta T_f = 273.15 - 270.65 = 2.5 \text{ K}$

$$\Delta T_f = K_f m \Rightarrow 2.5 = 1.86 \times \frac{n}{0.1}$$

$$\Rightarrow n = 0.1344 \text{ moles}$$


$$\Rightarrow w = 0.1344 \times 32 = 4.3 \text{ g}$$

$$\text{Volume} = \frac{4.3}{0.792} = 5.43 \text{ ml} = 543 \times 10^{-2} \text{ ml}$$




The ratio of number of oxygen atoms to bromine atoms in the product Q is _____ × 10⁻¹.


Ans. (15)



85. Number of carbocation from the following that are **not** stabilized by hyperconjugation is.....



Ans. (5)



86. For the reaction at 298 K, 2A + B → C. $\Delta H = 400 \text{ kJ mol}^{-1}$ and $\Delta S = 0.2 \text{ kJ mol}^{-1} \text{ K}^{-1}$. The reaction will become spontaneous above _____ K.

Ans. (2000)

Sol. $\Delta G = 0$

$$T = \frac{\Delta H}{\Delta S} = \frac{400}{0.2} = 2000 \text{ K}$$

87. Total number of species from the following with central atom utilising 2p² hybrid orbitals for bonding is.....

NH₃, SO₂, SiO₂, BeCl₂, C₂H₂, C₂H₄, BCl₃, HCHO, C₆H₆, BF₃, C₂H₄Cl₂

Ans. (6)

Sol. Central atom utilising sp² hybrid orbitals

SO₂, C₂H₄, BCl₃, HCHO, C₆H₆, BF₃



Download the new **ALLEN** app
& enroll for **Online Programs**

CLICK HERE TO
DOWNLOAD

88. Consider the two different first order reactions given below
 $A + B \rightarrow C$ (Reaction 1)
 $P \rightarrow Q$ (Reaction 2)
 The ratio of the half life of Reaction 1 : Reaction 2 is 5 : 2. If t_1 and t_2 represent the time taken to complete $\frac{2}{3}^{\text{rd}}$ and $\frac{4}{5}^{\text{th}}$ of Reaction 1 and Reaction 2, respectively, then the value of the ratio $t_1 : t_2$ is _____ $\times 10^{-1}$ (nearest integer).
 [Given: $\log_{10}(3) = 0.477$ and $\log_{10}(5) = 0.699$]

Ans. (17)

Sol. $\frac{(t_{1/2})_I}{(t_{1/2})_{II}} = \frac{K_2}{K_1} = \frac{5}{2}$
 $\therefore K_1 t_1 = \ln \frac{1}{1 - \frac{2}{3}} = \ln 3$
 $K_2 t_2 = \ln \frac{1}{1 - \frac{4}{5}} = \ln 5$
 $\Rightarrow \frac{K_1}{K_2} \times \frac{t_1}{t_2} = \frac{0.477}{0.699}$
 $\Rightarrow \frac{t_1}{t_2} = \frac{0.477}{0.699} \times \frac{5}{2} = 1.7 = 17 \times 10^{-1}$

89. For hydrogen atom, energy of an electron in first excited state is -3.4 eV, K.E. of the same electron of hydrogen atom is x eV. Value of x is _____ $\times 10^{-1}$ eV. (Nearest integer)

Ans. (34)

90. Among VO_2^+ , MnO_4^- and $\text{Cr}_2\text{O}_7^{2-}$, the spin-only magnetic moment value of the species with least oxidising ability is.....BM (Nearest integer).

(Given atomic member V = 23, Mn = 25, Cr = 24)

Ans. (0)

Sol. For 3d transition series;
 Oxidising power : $\text{V}^{+5} < \text{Cr}^{+6} < \text{Mn}^{+7}$
 $\text{V}^{+5} : [\text{Ar}] 4s^0 3d^0$
 Number of unpaired electron = 0

$\mu = 0$



Download the new ALLEN app & enroll for Online Programs

CLICK HERE TO DOWNLOAD